Roll No.	•••••
----------	-------

DD-766

M. A./M. Sc. (Fourth Semester)

EXAMINATION, 2020

MATHEMATICS

Paper Third (C)

[(Fuzzy Set Theory and Its Applications (II)]

Time: Three Hours

Maximum Marks: 80

Note: Attempt any *two* parts from each question. All questions carry equal marks.

Unit-I

- (a) Define logic, propositional logic. Write canonical form of modus ponens, modus tollens, hypothetical syllogism, unconditional and qualified proposition, conditional and unqualified proposition, conditional and qualified propositions.
 - (b) Give the steps of truth value restriction.

P. T. O.

(c) Let:

$$X = x_1, x_2, x_3$$

$$Y = y_1, y_2$$

$$Z = z_1, z_2$$
and
$$A = \left(\frac{.5}{x_1}, \frac{1}{x_2}, \frac{6}{x_3}\right)$$

$$B = \left\{\frac{1}{y_1}, \frac{.4}{y_2}\right\}, C = \left\{\frac{.2}{z_1}, \frac{1}{z_2}\right\}$$
for J $a, b = \begin{cases} 1 & \text{if } a \le b \\ b & \text{if } a > b \end{cases}$

then find:

$$R_3 x, z = \underset{y \in Y}{\operatorname{Sup \, min}} R_1 x, y, R_2 y, z$$

Unit—II

- 2. (a) Draw architecture of expert system.
 - (b) Show that:

$$J a,b = f^{-1}$$

$$f 1 - f a + f b ,$$

where $f: 0,1 \rightarrow 0,\infty$, f 0 = 0

is an increasing function, is a fuzzy implication.

(c) If:

$$A_{1} = \left(\frac{1}{x_{1}}, \frac{.9}{x_{2}}, \frac{.1}{x_{3}}\right)$$

$$A_{2} = \left(\frac{.9}{x_{1}}, \frac{1}{x_{2}}, \frac{.2}{x_{3}}\right)$$

$$B_{1} = \left(\frac{1}{y_{1}}, \frac{.2}{y_{2}}\right)$$

$$B_{2} = \left(\frac{.2}{y_{1}}, \frac{.9}{y_{2}}\right)$$

$$A_{3} = \frac{.8}{x_{1}}, \frac{.9}{x_{2}}, \frac{.1}{x_{2}}$$

Find B₃ by method of interpolation.

Unit-III

- (a) Discuss the main issues involved in the design of a fuzzy controller for stabilizing an inverted pendulum.
 - (b) Write a short note on fuzzification of classical dynamic systems.
 - (c) Write assumptions in a fuzzy control system design.

Unit-IV


- 4. (a) What do you mean by defuzzification? Write a brief account of centre of sums method.
 - (b) Aggregate graphically the fuzzy sets:

$$A_1 = \frac{0}{0}, \frac{.3}{1}, \frac{.3}{2}, \frac{.3}{3}, \frac{.3}{4}, \frac{0}{5}$$

$$A_2 = \frac{0}{3}, \frac{.5}{4}, \frac{.5}{5}, \frac{.5}{6}, \frac{0}{7}$$

$$A_3 = \frac{0}{5}, \frac{1}{6}, \frac{1}{7}, \frac{0}{8}$$

(c) Find x^* by method of centroid method for the figure :



- 5. (a) If $^{0+}A = 0,4$, $^{1}A=1,3$ and B, C are symmetric triangular fuzzy numbers with centres $C_B=4$, $C_C=5$ and spreads $S_B=S_C=2$. Rank these fuzzy numbers with Hamming distance method.
 - (b) Explain the method of symmetric fuzzy linear programming method.
 - (c) Explain the method of proposed by Shimura to construct an odering of all given alternatives.

DD-766 3,900